

A Comment on the Zero Temperature Chiral Phase Transition in $SU(N)$ Gauge Theories

R. Sekhar Chivukula*

Department of Physics, Boston University,
590 Commonwealth Ave., Boston MA 02215

December 5, 1996

Abstract

Recently Appelquist, Terning, and Wijewardhana investigated the zero temperature chiral phase transition in $SU(N)$ gauge theory as the number of fermions N_f is varied. They argued that there is a critical number of fermions N_f^c , above which there is no chiral symmetry breaking and below which chiral symmetry breaking and confinement set in. They further argued that that the transition is not second order even though the order parameter for chiral symmetry breaking vanishes continuously as N_f approaches N_f^c on the broken side. In this note I propose a simple physical picture for the spectrum of states as N_f approaches N_f^c from below (*i.e.* on the broken side) and argue that this picture predicts very different and *non-universal* behavior than is the case in an ordinary second order phase transition. In this way the transition can be *continuous* without behaving conventionally. I further argue that this feature results from the (presumed) existence of an infrared Banks-Zaks fixed point of the gauge coupling in the neighborhood of the chiral transition and therefore depends on the long-distance nature of the non-abelian gauge force.

*e-mail addresses: sekhar@bu.edu

Recently Appelquist, Terning, and Wijewardhana [1] have investigated the zero temperature chiral phase transition in $SU(N)$ gauge theory as the number of massless Dirac fermions N_f is varied. To second order, the beta-function of such a theory is given by

$$\mu \frac{\partial}{\partial \mu} \alpha(\mu) = \beta(\alpha) \equiv -b \alpha^2(\mu) - c \alpha^3(\mu) - d \alpha^4(\mu) - \dots, \quad (1.1)$$

where

$$b = \frac{1}{6\pi} (11N - 2N_f) \quad (1.2)$$

$$c = \frac{1}{24\pi^2} \left(34N^2 - 10NN_f - 3 \frac{N^2 - 1}{N} N_f \right). \quad (1.3)$$

For a small number of flavors the theory is asymptotically free and one expects QCD-like behavior, with confinement and with the chiral $SU(N_f)_L \times SU(N_f)_R$ symmetry broken to its vectorial subgroup. If the number of flavors is large enough (in perturbation theory, greater than $11N/2$) asymptotic freedom (and hence chiral symmetry breaking and confinement) is lost. For a range of N_f less than $11N/2$ the first term in the β -function is negative and the theory is asymptotically free, but there appears (in perturbation theory) to be a nontrivial infrared (Banks-Zaks) fixed point [2] because the second term in the β -function is *positive*. For N_f just slightly less than $11N/2$, this fixed point α^* is at weak coupling and the analysis is self-consistent. As N_f is lowered further, the fixed point moves to *larger* coupling.

In vector-like gauge theories an analysis of the gap equation [3, 4] suggests that, in a theory with an approximately constant coupling, chiral symmetry breaking occurs only if the coupling α exceeds a critical value

$$\alpha_c \equiv \frac{\pi}{3C_2(R)} = \frac{2\pi N}{3(N^2 - 1)}. \quad (1.4)$$

The authors of [1] suggested that the Banks-Zaks fixed point [2] persists to large coupling, and that the chiral-symmetry breaking transition in N_f should be associated with the point where $\alpha^* = \alpha_c$. They thereby estimated that

$$N_f^c = N \left(\frac{100N^2 - 66}{25N^2 - 15} \right). \quad (1.5)$$

Furthermore, Appelquist, Terning and Wijewardhana suggested that the nature of this phase transition is peculiar. Based on a gap-equation analysis [5] of the dynamical mass $\Sigma(p)$ of the fermions in the broken phase, they argued that the order parameter for chiral symmetry breaking (which is proportional to $\Sigma(0)$)

$$\Sigma(0) \approx \Lambda \exp \left(\frac{-\pi}{\sqrt{\frac{\alpha_*}{\alpha_c} - 1}} \right). \quad (1.6)$$




Figure 1: Spectrum of bosonic excitations in NJL model for couplings close to κ_c . Chiral symmetry is broken for $\kappa > \kappa_c$, and preserved for $\kappa < \kappa_c$. The σ and π in the broken phase combine to be the scalar ϕ multiplet in the unbroken phase. The vector and axial vector resonances form one chiral multiplet, and are therefore degenerate, in the unbroken phase.

goes to zero continuously¹ as $N_f \rightarrow (N_f^c)^-$ (and $\alpha^* \rightarrow \alpha_c^+$). Here the high-energy scale Λ represents the scale at which the coupling is far enough from the fixed point value to begin to run [1].

On the other hand, based of an analysis of the Bethe-Salpeter equation for the quark-antiquark scattering amplitude, Appelquist, Terning, and Wijewardhana argue that in the symmetric phase (close to the transition, when N_f is just above N_f^c) there are no light scalar resonances. Indeed, since the theory with $N_f \gtrsim N_f^c$ is presumed to be in a conformally-invariant “non-abelian coulomb” phase [2], it cannot have *any* isolated states. They then concluded that the phase transition is not second order *even though the order parameter changes continuously*.

In this note I propose a simple physical picture for the spectrum of states as N_f approaches N_f^c from below (*i.e.* on the broken side) and argue that this picture predicts very different and *non-universal* behavior than is the case in an ordinary second order phase transition. In this way the transition can be *continuous* without behaving conventionally [7]. I further argue that this feature results from the (presumed) existence of an infrared Banks-Zaks fixed point of the gauge coupling in the neighborhood of the chiral transition and therefore depends on the long-distance nature of the non-abelian gauge force.

Before moving to the case of a long-range force, let us review the familiar be-

¹ For momenta below $\Sigma(0)$, the fermions can be integrated out and no longer contribute to the β -function. Therefore, strictly speaking, for any $N_f < N_f^c$ there is no fixed-point. Nonetheless, for N_f close to N_f^c the coupling remains close to α^* for a large range of momenta, in a manner reminiscent of “walking technicolor” [6].




Figure 2: Spectrum of bosonic excitations in $SU(N)$ gauge theory for N_f close to N_f^c . For $N_f < N_f^c$, chiral symmetry is broken. For $N_f \gtrsim N_f^c$, the theory is assumed to be in a conformally-invariant “non-abelian coulomb” (NAC) phase [2] which has no isolated single-particle states.

havior of the chiral phase transition in a model with short-range interactions, the NJL model [8]. Here the fundamental interactions are modeled by chirally-invariant local four-fermion operators

$$\mathcal{L} = -\frac{4\pi\kappa}{\Lambda^2} \left[\bar{\psi} \gamma_\mu \frac{\lambda^a}{2} \psi \right]^2, \quad (1.7)$$

where the $\lambda^a/2$ are the generators of $SU(N)$ “color” and the ψ are the N_f flavors of fermions. This interaction is attractive in the chiral symmetry breaking channel, but is suppressed by a (large) energy scale Λ . In the limit where the NJL coupling κ is small, chiral symmetry remains unbroken. When the coupling is large, the chiral symmetry breaking scale (as characterized by the value of the F -constant — the analog of f_π in QCD — or by Σ , the momentum-independent dynamical mass of the fermion) is of order Λ . There is a critical value κ_c , estimated to be $\pi/3$ in the large- N limit, below which chiral symmetry is unbroken and above which it is broken. If the transition between these two regimes is *smooth*, as it is in the gap-equation in the fermion-bubble approximation, the dynamical mass of the fermion goes smoothly to zero as $\kappa \rightarrow (\kappa_c)^+$, and remains identically zero for $\kappa \leq \kappa_c$.

In the case of the Nambu-Jona-Lasinio model, the spectrum of bosonic excitations [10] is shown in Figure 1. Note that, aside from the light scalar-multiplet ϕ on the unbroken side and the σ and Goldstone bosons π on the broken side, *all* other “excitations” have a mass of order Λ ! The reason for this is that the intrinsic physical scale of the interactions is *always* of order Λ . Near $\kappa = \kappa_c$, the scalar states are *anomalously light* due to the “fine-tuning” [9] of the NJL interaction.

This picture should be contrasted with the analogous change of the spectrum

of particles near the chiral phase transition in non-abelian gauge theory as $N_f \rightarrow (N_f^c)^-$. Because of the infrared fixed-point, the high-energy scale Λ (see eqn. 1.6) is no longer relevant. In this case the *only* relevant dynamical scale is the magnitude of dynamical mass $\Sigma(0)$. All other scales, the F -constant, the confinement scale, *etc.* are of the same order of magnitude [1]. Therefore, if $\Sigma(0) \rightarrow 0$ continuously as $N_f \rightarrow (N_f^c)^-$, we expect the spectrum of bosonic excitations to be as shown in Figure 2. Note that as $N_f \rightarrow (N_f^c)^-$, the entire spectrum collapses to zero mass. So long as the Banks-Zaks fixed point persists² in the non-abelian gauge theory, all high-energy scales are irrelevant. Therefore, unlike the NJL model, if $\Sigma(0) \rightarrow 0$ the mass of all excitations must also tend to zero.

What are the implications of this behavior? In the case of the NJL model near the critical coupling, it is appropriate to “integrate-out” all higher-mass states and the critical behavior of the theory is determined entirely by the infrared behavior of the corresponding scalar field theory³. We therefore expect similar behavior in *any* $SU(N_f) \times SU(N_f)$ chirally-invariant four-dimensional field theory with *short-range interactions* near the phase boundary: *i.e.* we expect the behavior to be *universal*.

In contrast, in the case of non-abelian gauge theory with N_f just below N_f^c , we *cannot* reduce the theory to an effective low-energy scalar theory. In principle *all* “higher” resonances contribute to any correlation function and the behavior is *non-universal*. In this case it is plausible that the order parameter changes continuously as $N_f \rightarrow (N_f^c)^-$, even though the theory with $N_f \gtrsim N_f^c$ has no light scalar states.

Finally, I note that arguments similar to those presented here may cast light on the discrepancies between the gap-equation [14, 15] and the effective scalar field-theory [16] analyses of the chiral phase transition in QED3.

In this note I have proposed a simple physical picture for the spectrum of states in $SU(N)$ gauge theory as the number of flavors N_f approaches the critical number N_f^c from below (*i.e.* from the broken side). This picture predicts very different and *non-universal* behavior than is the case in a conventional second order phase transition. I have further argued that this feature results from the (presumed) existence of a Banks-Zaks fixed point of the gauge coupling in the neighborhood of the chiral transition and therefore depends on the long-distance nature of the non-abelian gauge force.

Acknowledgments

I thank Koichi Yamawaki and the organizers of the 1996 International Workshop on Perspectives of Strong Coupling Gauge Theories (SCGT 96) held in Nagoya, Japan from 13-16 November, 1996 for holding a stimulating conference where this

²More properly, since we are in the broken phase where no true fixed-point exists, so long as the coupling remains close to α^* over a large range of momenta.

³In some cases the resulting scalar theory cannot have a second order transition, but will have a fluctuation-induced first-order transition instead [11]. If the transition is driven to be first-order, all relevant dimensional quantities will be [12, 13] of order Λ .

work was begun. I also thank Nick Evans, Stephen Selipsky, Rohana Wijewardhana, Elizabeth Simmons and John Terning for discussions and for comments on the manuscript. *This work was supported in part by the Department of Energy under grant DE-FG02-91ER40676.*

References

- [1] T. Appelquist, J. Terning, and L. C. R. Wijewardhana, *Phys. Rev. Lett.* **77** (1996) 1214.
- [2] T. Banks and A. Zaks, *Nucl. Phys.* **B196** (1982) 189.
- [3] T. Maskawa and H. Nakajima, *Prog Theor. Phys* **52** (1974) 1326; **54** (1976) 860; R. Fukuda and T. Kugo *Nucl. Phys.* **B117** (1976) 250; P.I. Fomin and V.A. Miransky, *Phys. Lett.* **B64** (1976) 166.
- [4] J.M. Cornwall, R. Jackiw, and E. Tomboulis, *Phys. Rev.* **D10** (1974) 2428.
- [5] V.A. Miransky *Il Nuovo Cimento* **90A** (1985) 149; V.A. Miransky and P.I. Fomin, *Sov. J. Part. Nucl.* **16** (1985) 203.
- [6] B. Holdom, *Phys. Rev* **D24** (1981) 1441; B. Holdom, *Phys. Lett.* **B150** (1985) 301; K. Yamawaki, M. Bando, and K. Matumoto, *Phys. Rev. Lett.* **56** (1986) 1335; T. Appelquist, D. Karabali, and L.C.R. Wijewardhana, *Phys. Rev. Lett.* **57** (1986) 957; T. Appelquist and L.C.R. Wijewardhana, *Phys. Rev* **D35** (1987) 774; T. Appelquist and L.C.R. Wijewardhana, *Phys. Rev* **D36** (1987) 568.
- [7] For an alternative, and possibly complementary, view of the physics of this transition, see V. A. Miransky and K. Yamawaki, hep-th/9611142.
- [8] Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122**, 345 (1961).
- [9] G. 't Hooft, in *Recent Developments in Gauge Theories*, G. 't Hooft, *et. al.*, eds., Plenum Press, New York NY 1980.
- [10] For the ρ -meson see, for example, V. Bernard and U. G. Meissner, *Nucl. Phys.* **A489** (1988) 647.
- [11] S. Coleman and E. Weinberg, *Phys. Rev.* **D7** (1973) 1888.
- [12] R. S. Chivukula, M. Golden, and E. H. Simmons, *Phys. Rev. Lett.* **70** (1993) 1587.
- [13] W. A. Bardeen, C. T. Hill, D. Jungnickel, *Phys. Rev.* **D49** (1994) 1437.
- [14] T. Appelquist, D. Nash, and L.C.R. Wijewardhana, *Phys. Rev. Lett.* **60**, 2575 (1988); D. Nash *Phys. Rev. Lett.* **62** (1989) 3024; T. Appelquist and D. Nash, *Phys. Rev. Lett.* **64**, 721 (1990).
- [15] T. Appelquist, J. Terning, and L.C.R. Wijewardhana, *Phys. Rev. Lett.* **75** (1995) 2081.
- [16] R. D. Pisarski, *Phys.Rev.* **D44** (1991) 1866.